

Uses of bacteriophages

Name: Chris Wong
Supervisor: Prof. Margaret Ip
Department of Microbiology
Date: 3-12-2013


Discovery of bacteriophages

- World War I
- Viruses "eaters of micro-organism" discovered by Felix d'Herelle
- Isolation of filterable entities able to lyse bacterial culture
- Small cleared zones on plate called "plaques"
- Treat World War I soldiers affected by *Shigella* dysentery
- Have been used clinically in Eastern Europe and the former Soviet Union since 1919
- Polish and Soviets administered phage orally, topically or systemically to treat antibiotic resistant pathogens, including *Staphylococcus*, *Streptococcus*, *Klebsiella*, *Escherichia*, *Proteus*, *Pseudomonas*, *Shigella* and *Salmonella*
- Success rate : 80-95% of phage therapy
- Not taken seriously in the West due to discovery of antibiotics
- Emergence of multi-drug-resistant pathogen, growing interest in phage therapy




Bacteriophages(phages)

- Most abundant 'life' forms on Earth
- Water, soil and air
- Estimated 10^{32} bacteriophages in total on the planet
- Specifically target bacteria, not mammalian cells
- Each phage only attacks one species
- Head: Icosahedral shape capsid, RNA/DNA
- Tail: usually 6 fibers with receptors at tips, recognition of attachment sites on bacterial cell surface e.g. teichoic acid, oligosaccharide, peptidoglycan
- Two types of life cycles: lytic (virulent phage) or lysogenic (temperate phage) cycle



Lytic cycle and lysogenic cycle




Potential use of bacteriophages in treatment of bacterial infections

- Emergency of multi-drug-resistant strains of pathogens
- New antibiotics development rate slow
- Cannot replace old drugs
- Alternative strategies for treatment of antibiotic-resistant pathogens
- Lytic phages preferred
- lysogenic phages, no host death, transfer undesirable genes (e.g. toxin genes/antibiotic resistance genes) through transduction
- Advantages over antibiotic:
 - Narrower host range
 - No side effect
 - Replicate exponentially, a self-regulating tool
 - None can replicate in eukaryotes

Isolation of bacteriophages

- Sewage water
- Bacterial culture in broth
- Inoculate the sewage water
- (specific phage binds to and replicates in host cells, lyses the bacteria eventually)
- Centrifuge to remove all cell debris
- Filter supernatant with 0.22 micron filter
- Spread O/N culture on agar plate (confluent lawn of bacteria)
- Add filtrate to center of plate
- A plaque (clear zone) will be observed where phage infect and lyse the bacteria
- phage capable of infecting the host strain has been isolated



Phage therapy: treatment of diabetic foot infections

- Wound infections leading to foot and leg amputation: 30-50% diabetic patients
- 43% MRSA infections
- Potential use of lytic phages for treating MRSA infections in diabetic patients
- Phage MR-10 (broad host range)
- Localized foot infection caused by MRSA in diabetic mice evaluated

Phage therapy: treatment of diabetic foot infections

- BALB/c mice injected with alloxan monohydrate
- toxic glucose analogue (150mg/kg body weight)
- Selectively damage beta-cells(insulin-producing cells) in the pancreas
- Lytic Phage MR-10
- Linezolid
- Combination therapy

Phage therapy: treatment of diabetic foot infections




Lesion scoring:



- 0- No redness, no oedema with thickness of hindpaw in the range of 0.14–0.19 mm.
- 1- Slight redness with slight oedema with thickness of hindpaw in the range of 0.20–0.25 mm.
- 2- Visible redness and moderate oedema with thickness of hindpaw in the range of 0.26–0.30 mm.
- 3- Severe redness and visible oedema with thickness of hindpaw in the range of 0.31–0.36 mm.
- 4- Severe redness and pronounced visible oedema with thickness of hindpaw in the range of 0.37–0.42 mm.

Phage therapy: treatment of diabetic foot infections



Phage therapy: treatment of diabetic foot infections

Days	Untreated Control		Phage MR-10 Treated		LZD treated (25 mg/kg/per oral)		Phage MR-10 + LZD treated	
	Lesion score	Oedema (mm)	Lesion score	Oedema (mm)	Lesion score	Oedema (mm)	Lesion score	Oedema (mm)
Day 0	0	0.16±0.01	0	0.17±0.01	0	0.17±0.02	0	0.18±0.02
Day 1	1	0.25±0.02	1	0.22±0.02	1	0.24±0.02	1	0.22±0.02
Day 3	3	0.35±0.02	1	0.28±0.01	2	0.3±0.02	2	0.25±0.02
Day 5	4	0.40±0.01	2	0.26±0.02	1	0.25±0.04	1	0.21±0.02
Day 7	3	0.32±0.02	1	0.2±0.01	0	0.18±0.01	0	0.17±0.01
Day 9	1	0.21±0.01	0	0.18±0.02	0	0.18±0.02	0	0.16±0.01
Day 12	0	0.17±0.01	0	0.17±0.01	0	0.16±0.02	0	0.17±0.01

Data is mean±S.D of a minimum of four independent values.
doi:10.1371/journal.pone.0056022.t002

Phage therapy: treatment of diabetic foot infections

- Similar efficacy to linezolid
- Combination therapy more effective in controlling entire process of hindpaw infection in diabetic mice compared to antibiotic/phage alone
- Linezolid resistant strains emergence
- Decrease development of resistant strain
- Effective strategy for patients not respond to conventional antibiotic therapy

Phage Therapy: rescue mice with VRE bacteremia

- GI tract colonization of vancomycin-resistant *Enterococcus faecium*(VRE)
- Predispose individual to VRE bacteremia, esp. immunocompromised patients
- Clinical isolates of VRE used to induce bacteremia in mice by intraperitoneal (i.p.) injection of 10^9 CFU
- Fatal within 48 h
- *Enterococcus* phages ENB6 isolated from raw sewage at a sewage treatment plant
- Single i.p. injection of 3×10^8 PFU of lytic phage administered 45 min after infection
- Rescue 100% of mice
- Even treatment delayed to the point where all mice are moribund
- ~50% mice rescued by a single injection of phages

Phage Therapy for infections in cancer patients

- Performed in 20 cancer patients (ages 1–66 years, 17 with solid tumors, 3 with hematological malignancies)
- All patients had concurrent bacterial infections (*Staphylococcus aureus*, 8 patients; *Pseudomonas aeruginosa*, 9; *Klebsiella pneumoniae*, 6; *Klebsiella oxytoca*, 2; *Escherichia coli*, 6)
- Homogenous (12 patients) and mixed (8 patients) infections
- All patients previously treated with antibiotics without apparent response
- In all patients, bacteriophages were administered orally 3 times daily.
- Eight of them had localized bacteriophage treatment in addition to the oral protocol.
- No side effects of the therapy were observed
- Cure of infection achieved in all cases (cessation of suppuration, closure of wounds, eradication of pneumonia, etc.)

Safety of phage therapy in human

- Human volunteers receiving *Escherichia coli* phage T4 orally: a safety test of phage therapy
 - T4 dose (10^3 PFU/ml), a higher phage dose (10^5 PFU/ml)
 - No adverse events related to phage application were reported
 - neither T4 phage nor T4-specific antibodies were observed in the serum of the subjects at the end of the study
- Bacteriophage therapy of venous leg ulcers in humans: results of a phase I safety trial
 - 42 patients with chronic venous leg ulcers, 39 patients completed the trial
 - treated for 12 weeks with either a saline control or bacteriophages targeted against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Escherichia coli*
 - No increase in adverse reaction associated with phage cocktail application

Commercial phage products

- JSC Biochimpharm, Biopharm L Limited
 - Republic of Georgia
 - Various phage lysates mixed for intestinal problems e.g. dysentery, salmonellosis and colitis
 - Patented and licensed liquid and tablet phage product
- EBI Food Safety
 - Netherlands
 - LISTEX P100™ a cocktail of phage against *Listeria*
- Phico Therapeutics (UK), Viridax(USA)
 - Anti-MRSA phage product
 - Pre-clinical
- BioControl
 - Uk
 - Phage product for *Pseudomonas* infections of ear
 - Phase II clinical trial completed

Enzybiotics as antimicrobials



- Bacteriophage-derived endolysins
- Released from host cell by 2 phage genome-encoded protein:
- Holins disrupt cell membrane
- Endolysins digest cell wall, peptidoglycan hydrolase
- Contain two functional domains:
- cell wall binding module(CWBM), bind to species-specific carbohydrate epitope in cell wall
- catalytic domain, cleave peptidoglycan bonding
- Lyse Gram +ve bacteria when applied exogenously
- *Staphylococcus aureus* Phage P68 *lys16* gene, cell lysis in clinical isolates of *S. aureus*
- To date, no reports of resistant strain against phage endolysins


Enzybiotics as antimicrobials

- Novel Bacteriophage lysin, broad lytic activity
- Phage lysin (PluSs2) derived from *Streptococcus suis* phage
- Activity against MRSA, VISA, *S. epidermidis*, *Listeria*, GBS, GES, GGS, *S. pyogenes*, *S. pneumoniae*
- Protect against mixed infection
- 128 g/ml *in vitro* reduced MRSA and *S. pyogenes* growth by 5 logs and 3 logs within 1 h respectively
- A single, 2-mg dose of PlySs2 protected 92% (22/24) of the mice in a bacteremia model of mixed MRSA and *S. pyogenes* infection
- Effective therapeutic


Phages as biocontrol agents

- Reduce biofilm formation on indwelling catheters
- 1/3-1/2 nosocomial endocarditis cases due to infected intravascular catheters in US
- ~250,000 cases of catheter-related bloodstream infections, mortality 12-25%
- *S. aureus*, *S. epidermidis*, forming biofilm
- Current treatment: antimicrobial coating of catheters, disinfection of insertion site
- Biofilm, antimicrobial resistant strain
- A novel approach using a phage pretreatment step
- biofilm formation by *S. epidermidis* after 24 h on catheter significantly reduced in the presence of phage 456





Catheter
after biofilm
formation by
P. aeruginosa
for 24 h



Catheter
pretreated
with *P.*
aeruginosa
phage M4
and exposed
for 24 h to *P.*
aeruginosa

Phages as biocontrol agents

- Application of phages to control bacteria on food
- Food preservation and safety
- Food-borne illness by *Salmonella*
- Phage cocktail composed of 3 lytic phages (UAB_Phi 20, UAB_Phi78, and UAB_Phi87)
- Food (pig skin, chicken breasts and packaged lettuce) experimentally contaminated with *Salmonella*
- Significant bacterial reduction treated with phage cocktail

Phages as biocontrol agents

- US Food and Drug Administration, 2006
- Approved use of phage to be added to meat and poultry products
- Mixture 6 Purified phages allowed to be used on read-to-eat (RTE) meat and poultry products
- As antimicrobial agent against *Listeria monocytogenes*
- Effective against 170 strains
- Directly consumed without additional cooking, capable multiply at low temperatures
- Spray application to RTE food surface e.g. lunch meats and hot dogs prior to packaging
- Commercially available phage preparations for food safety applications Listex™ P100 (left) and ListShield™ (right).




Phages as biocontrol agents

- Phage VPp1 against *Vibrio parahaemolyticus*
- Closely associated with oysters normally consumed raw or lightly cooked
- Phage FAHc1 against *E. coli* O157 on beef
- Listex P100 to control *L. monocytogenes* on fresh-cut fruits and fruit juices
- *Lactococcus garvieae*, opportunistic pathogens, infectious diseases in fish
- Phage PLgY in aquaculture


Genetically engineered phages as molecular tools

- Identification of bacterial targets
- Fluorescent labeling of phage nucleic acid or phage head or tail component
- Visual tag outline and identify bacterial cell for which the phage is host-specific




Genetically engineered phages as molecular tools


- Identification of bacterial targets
- Manipulate phage's genome to carry a reporter gene e.g. green fluorescent protein (GFP)
- Expressed when delivered to bacterial host
- Visualized by fluorescent signal



Genetically engineered phages as molecular tools


- Diagnosis of *Mycobacterium tuberculosis*
- Slow-growing, culturing from sputum samples requires up to 6 weeks before identification
- Commercial kit *FASTPlaqueTB*TM
- Sputum sample mixed with phage suspension for attachment to any MTB cells
- Unbound phages killed by virucide, derived from pomegranate extract
- Those attached not killed
- Mixed with a fast-growing non-pathogenic species of *Mycobacterium* which the phage also capable to infect
- Spread on agar
- Fast-grower produces a confluent lawn
- Plaques due to presence of MTB cells in sputum sample that infected prior to virucide treatment
- Plaque indicate patient suffering from TB





Genetically engineered phages as molecular tools

- RNA interference (RNAi)
- Gene silencing
- Delivery vehicles



Conclusion

- Phage therapy
 - Lytic phages against pathogens
 - Effective, safe and low cost
 - alternative strategy for multi-drug-resistant strains
- Enzybiotics as antimicrobials
 - Endolysin for Gram +ve bacterial infection
- Phages as biocontrol agents
 - Biofilm on indwelling catheters
 - Food preservation
- Phages as molecular tools
 - Bacterial identification
 - Nucleic acid delivery

References

- Alisky J, Iczkowski K, Rapoport A, Troitsky N (1998) Bacteriophages Show Promise as Antimicrobial Agents. *Journal of Infection* 36: 5-15.
- Hanlon GW (2007) Bacteriophages: an appraisal of their role in the treatment of bacterial infections. *International Journal of Antimicrobial Agents* 30(2):118-128
- Brussow H, Kutter E. Phage ecology. 2005 . In: Kutter E, Sulakvelidze A, editors. *Bacteriophages: biology and applications*. Boca Raton, FL: CRC Press: 129–63.
- Kutateladze M, Adamia R (2010) Bacteriophages as potential new therapeutics to replace or supplement antibiotics. *Trends in Biotechnology* 28(12): 591-595
- Hodgson K (2013) Bacteriophage therapy. *Microbiology Australia*. 28-31.
- Cervený KE, DePaola A, Duckworth DH, Gulig PA (2002) Phage therapy of local and systemic disease caused by *Vibrio vulnificus* in iron - dextran -treated mice. *Infection and Immunology* 70(11): 6251–6262
- Bruttin A . Brussow H (2005) Human Volunteers Receiving *Escherichia coli* Phage T4 Orally: a Safety Test of Phage Therapy. *Antimicrob Agents Chemother*. 49(7): 2874–2878
- Rhoads DD, WolcottRD, Kuskowski Ma, WolcottBM, Ward LS, Sulakvelidze A (2009) Bacteriophage therapy of venous leg ulcers in humans: results of a phase I safety trial. *J Wound Care*18(6):237-8, 240-3
- Chanishvili N (2012) Phage Therapy—History from Twort and d'Herelle Through Soviet Experience to Current Approaches. *Advances in Virus Research* 83:3-40
- Chhibber S, Kaur T, Kaur S (2013) Co-Therapy Using Lytic Bacteriophage and Linezolid: Effective Treatment in Eliminating Methicillin Resistant *Staphylococcus aureus* (MRSA) from Diabetic Foot Infections. *PLoS ONE* 8(2): e56022.
- Biswas B, Adhya S, Washart P, Paul B, Trostel AN, Powell B, Carlton R, Merrill CR (2002) Bacteriophage Therapy Rescues Mice Bacteremic from a Clinical Isolate of Vancomycin-Resistant *Enterococcus faecium*. *Infection and Immunity* 70(1): 204-210
- Weber-Dabrowska B, Górski MM (2000) Bacteriophage therapy for infections in cancer patients. *Clinical and Applied Immunology Reviews* 1: 131–134
- Housby JN, Mann NH (2009) Phage therapy *Drug Discovery Today* 14: 536-540
- Nelson DC, Schmelcher M, Rodriguez-Rubio L, Klumpp J, Pritchard DG, Dong S, Donovan DM (2012) Endolysins as antimicrobials. *Advances in Virus Research* 83:299-365
- Takáč M, Witte A, Bläsi U (2005) Functional analysis of the lysis genes of *Staphylococcus aureus* phage P68 in *Escherichia coli*. *Microbiology* 151(Pt7): 2331-42
- Hermoso JA, Garcia JL, Garcia P (2007) Taking aim on bacterial pathogens: from phage therapy to enzybiotics. *Current Opinion in Microbiology* 10:461–472
- Gilmer DB, Schmitz JE, Euler CW, Fischetti VA (2013) Novel Bacteriophage Lysin with Broad Lytic Activity Protects against Mixed Infection by *Streptococcus pyogenes* and Methicillin-Resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy*. 57(6): 2743-2750
- Curtin JJ, Donlan RM (2006) Using Bacteriophages To Reduce Formation of Catheter-Associated Biofilms by *Staphylococcus epidermidis*. *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY* 50(4): 1268-1275
- Fu W, Forster T, Mayer O, Curtin JJ< Lehman SM, Donlan (2009) Bacteriophage Cocktail for the Prevention of Biofilm Formation by *Pseudomonas aeruginosa* on Catheters in an *In Vitro* Model System. *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY* 54(1): 397-404.
- Spricigo DA, Bardina C, Cortés P, Llagostera M (2013) Use of a bacteriophage cocktail to control Salmonella in food and the food industry. *Int J Food Microbiol*. 165(2): 169-74
- Peek R, Reddy KR (2006) FDA Approves Use of Bacteriophages to be Added to Meat and Poultry Products. *GASTROENTEROLOGY* 131:1370–1372
- Sulakvelidze A (2013) Using lytic bacteriophages to eliminate or significantly reduce contamination of food by foodborne bacterial pathogens. *J Sci Food Agric* 93: 3137–3146
- Rong R, Lin H, Wang J, Khan MN, Li M, Reductions of *Vibrio parahaemolyticus* in oysters after bacteriophage application during depuration, *Aquaculture* doi:10.1016/j.aquaculture.2013.09.028
- Hudson JA, Billington C, Cornelius AJ, Wilson T, Premaratne SA, King NJ (2013) Use of a bacteriophage to inactivate *Escherichia coli* O157:H7 on beef. *Food Microbiology* 36:14-21
- Oliveira M, Viñas I, Colàs P, Anguera M, Usall J, Abadías M (2013) Effectiveness of a bacteriophage in reducing *Listeria monocytogenes* on fresh-cut fruits and fruit juices. *Food Microbiology* 38:137-142
- Nakai T, Park SC (2001) Bacteriophage therapy of infectious diseases in aquaculture.. *Research in Microbiology* 153 : 13–18
- Smartt AE, Xu T, Jegier P, Carswell JJ, Blount SA, Steven Ripp GSS (2012) Pathogen detection using engineered bacteriophages *Anal Bioanal Chem* 402:3127–3146
- Hanlon, G. W. (2013) Applications of Bacteriophage Technology, in Russell, Hugo & Ayliffe's: Principles and Practice of Disinfection, Preservation and Sterilization, 5th Edition (eds A. P. Fraise, J.-Y. Maillard and S. A. Sattar), Wiley-Blackwell, Oxford, UK. doi: 10.1002/9781118425831.ch22b
- Albert H, Trollip A, Seaman T, Mole RJ (2004) Simple, phage-based (FASTPlaque) technology to determine rifampicin resistance of *Mycobacterium tuberculosis* directly from sputum **INT J TUBERC LUNG DIS** 8(9):1114–1119
- Chen J, Xie J (2012) Progress on RNAi-based molecular medicines. *International Journal of Nanomedicine* 7: 3971–3980