

Submitted: 1/11/2021

Accepted: 4/10/2021

DOI: <https://doi.org/10.3201/eid2707.210054>

21-0054 Research

Tables: 0

Figures: 2

Appendix: 1

TOC title: Psychobehavioral Responses and Likelihood of Receiving COVID-19 Vaccines during the Pandemic, Hong Kong

Running head: Psychobehavioral Responses to COVID-19 Vaccines

Keywords: SARS-CoV-2, COVID-19, coronavirus, 2019 novel coronavirus disease, severe acute respiratory syndrome coronavirus 2, zoonoses, coronavirus disease, viruses, psychobehavioral, risk perception, preventive measures, vaccine uptake, longitudinal assessment, vaccines, Hong Kong

Suggested citation for this article: Kwok KO, Li KK, Tang A, Tsoi MTF, Chan EYY, Tang JWT, et al. Psychobehavioral responses and likelihood of receiving COVID-19 vaccines during the pandemic, Hong Kong. *Emerg Infect Dis.* 2021 Jul (*date cited*).

<https://doi.org/10.3201/eid2707.210054>

Psychobehavioral Responses and Likelihood of Receiving COVID-19 Vaccines during the Pandemic, Hong Kong

Kin On Kwok, Kin Kit Li, Arthur Tang, Margaret Ting Fong Tsoi, Emily Ying Yang Chan, Julian Wei Tze Tang, Angel Wong, Wan In Wei, Samuel Yeung Shan Wong

Author affiliations: The Chinese University of Hong Kong, Hong Kong (K.O. Kwok, M.T.F. Tsoi, E.Y.Y. Chan, A. Wong, W.I. Wei, S.Y.S. Wong); City University of Hong Kong, Hong Kong (K.K. Li); Sungkyunkwan University, Seoul, South Korea (A. Tang); University of Leicester, Leicester, United Kingdom (J.W.T. Tang)

To access temporal changes in psychobehavioral responses, we conducted a 5-round longitudinal population-based online survey (R1–R5) in Hong Kong during January–September 2020. Most respondents reported wearing masks (R1, 99.0% to R5, 99.8%) and performing hand hygiene (R1, 95.8% to R5, 97.7%). Perceived severity decreased significantly, from 97.4% (R1) to 77.2% (R5), and perceived self-susceptibility remained high (87.2%–92.8%). Female sex and anxiety were associated with greater adoption of social distancing. Intention to receive COVID-19 vaccines decreased significantly (R4, 48.7% to R5, 37.6%). Greater anxiety, confidence in vaccine, collective responsibility, and weaker complacency were associated with higher tendency to receive COVID-19 vaccines. Although its generalizability should be assumed with caution, this study helps to formulate health communication strategies and foretells the initial low uptake rate of COVID-19 vaccines, suggesting that social distancing should be maintained in the medium term. Future research should study COVID-19 transmission after a mix of social distancing and vaccination.

Since the World Health Organization declared the 2019 novel coronavirus disease (COVID-19) a pandemic on March 11, 2020 (1), COVID-19 has infiltrated every continent in the world (2). Hong Kong, a densely populated city located on the southern coast of China with ≈ 7.5 million citizens and a mean daily number of 12.5 social encounters per individual (3), recorded a laboratory-confirmed COVID-19 case in late January 2020 (4). Since then, Hong Kong has been adopting a suppress-and-lift strategy, under which lifting and reimposing of restrictions were based on epidemiologic thresholds (5). As of April 9, 2021, there have been 11,550 confirmed cases and 205 deaths (crude case fatality rate 1.8%) (6), and the fourth wave of COVID-19 epidemic in Hong Kong had just ended. With more available data on phase 3 clinical trials of candidate vaccines (7) and the authorization for emergency use, the COVID-19 vaccination program in Hong Kong began in late February 2021.

Surveillance of psychobehavioral responses during the epidemic plays an essential role to convey risk communication messages to the public. Previously we reported that the general population in Hong Kong had high levels of perceived risk and mild anxiety during the early phase of the COVID-19 epidemic; the prompt government interventions with widely adopted individual precautionary measures might be the determinants to slow down the transmission early in the outbreak (8). After our initial analysis, based on cross-sectional data (8), global researchers have applied similar protocols for the general public in Japan (9), Saudi Arabia (10), Italy (11) and the United Kingdom (12). However, the temporal variations of psychobehavioral responses have not been examined.

In addition to psychobehavioral responses, unique to COVID-19 is its unprecedented massive epidemic size compared with other recent outbreaks, such that vaccination becomes the exit strategy. However, despite vaccine availability, vaccine doubters may hamper the global effort against COVID-19 (13). Unraveling the reasons behind vaccine hesitancy and monitoring its trends over time will support the design of interventions to boost COVID-19 vaccine uptake.

We report a longitudinal analysis of 5 representative population-based surveys of adults in Hong Kong on their psychological, behavioral, and vaccine-related responses conducted during the first 2 waves of the COVID-19 epidemic. Our main objectives were tracking major psychobehavioral responses (including risk perception, psychological distress, and adoption of precautionary measures) over time and examining the determinants of the intention to receive the COVID-19 vaccine. As a complement, other psychobehavioral responses (such as knowledge about COVID-19) were also reported. These findings should have major implications for infection control policies and targeted mental health recommendations. Hong Kong has a high-income economy but had major social unrest in the prepandemic period in the population (14); thus the experience in Hong Kong may act as a reference for other similar populations to prepare for future epidemics.

Methods

Respondent Recruitment

A community cohort was established within 36 hours after the first COVID-19 confirmed case was identified in Hong Kong, and we followed it. District councilors shared an online survey link through channels in which they convey information to their targeted residents (8). We arranged 5 follow-up rounds (denoted as R1–R5) of online surveys during January–September 2020, each lasting for 3–6 weeks, as follows: R1, January 23–February 13; R2, March 6–April 14; R3, May 8–June 14; R4, July 15–August 7; and R5, August 8–September 15. Respondents were compensated with Hong Kong dollars (HKD)10 (R1), HKD20 (R2), and HKD30 (R3–R5) worth of cash vouchers.

Study Instrument

The study instrument was based on a questionnaire used during the initial phase of the COVID-19 epidemic in Hong Kong (8). In each round, we administered questions soliciting key information on demographics, health conditions, travel history, risk perceptions toward COVID-

19, anxiety and burnout, confidence in the local government and doctors, knowledge about COVID-19 transmission, and adoption and perceived effectiveness of preventive measures. In response to the funding commitments for COVID-19 vaccine development (15), starting with R4 we embedded vaccine-related questions along 2 dimensions: intention to receive COVID-19 vaccines when available and vaccine hesitancy. The definitions and response scales of these items are outlined in this section or are self-explanatory from the tables (Appendix, <https://wwwnc.cdc.gov/EID/article/27/7/21-0054-App1.pdf>).

Psychological Responses

Risk perceptions toward COVID-19 included perceived susceptibility (of oneself and one's family members), assuming no precautionary measure, and perceived severity. Starting with R3, we asked respondents to report their perceived susceptibility based on the situation during which they completed the survey (1, very likely; 5, very unlikely). In addition, respondents rated the level of disease severity of COVID-19 and other noncommunicable diseases and infectious diseases (1, very bad; 5, not bad at all).

We measured anxiety with the Chinese–Cantonese version of the Hospital, Anxiety and Depression Scale—Anxiety (HADS-A) (16). Respondents rated 7 statements on the basis of their feelings in the preceding 7 days on a 4-point scale, with a higher score indicating stronger anxiety (summative score: 0–7, normal; 8–10, borderline abnormal; 11–21, abnormal).

We measured burnout with a single-item measure: “Overall, based on your definition of burnout, how would you rate your level of burnout when facing COVID-19?” (1, “I have no symptoms of burnout”; 5, “I feel completely burned out and often wonder if I can go on facing COVID-19; I am at the point where I may need some changes or may need to seek some sort of help”). This single-item measure was refined from a nonproprietary validated burnout measure (17) to fit the current context and was asked starting with R3.

Behavioral Responses

Respondents rated the adoption (yes/no) (Appendix Table 1) and perceived effectiveness (1, very effective; 5, not very effective) (Appendix Table 2) of 17 precautionary measures against COVID-19. For the likelihood of COVID-19 vaccine uptake, respondents answered this question “If COVID-19 vaccines are available, how likely will you receive them?” (0, definitely not; 10, definitely). We measured vaccine hesitancy with a validated 15-item tool (18) with 3

items on each of 5 psychological antecedents (the 5Cs): confidence: trust in the safety and effectiveness of the vaccine, the system that delivers the vaccine, and the motivations of policymakers who decide on the need of the vaccine; complacency: not perceiving the disease as high risk and vaccination as necessary; constraints: barriers to vaccination; calculation: persons' engagement in extensive information searching; and collective responsibility: willingness to protect others through herd immunity. We used an average score for each antecedent. For collective responsibility, one reverse item, "When everyone is vaccinated, I don't have to get vaccinated, too," was excluded from the calculation. The vaccine-related items did not include any specific information about pharmaceutical companies or manufacturing countries.

Statistical Analysis

We summarized responses using descriptive statistics. To examine the overall linear trends in the responses and to account for the correlation diminishment resulting from responses from the same respondents across time, we adopted the generalized estimation equation framework featuring an autoregressive structure for within-subject correlations. To compare the proportion of responses in 2 time points, we used a partially overlapping samples z -test (19).

We adopted a multivariate regression model under the generalized estimation equation framework to identify the associated factors for higher tendency for social distancing (i.e., ≥ 5 social distancing measures) and higher uptake tendency for COVID-19 vaccines (i.e., ≥ 7 of 10 for the "likelihood of receiving COVID-19 vaccines" question). We reported adjusted odds ratios (aORs) and 95% confidence intervals. We specified a statistical significance of $p \leq 0.05$. We conducted the analysis in R version 4.0.3 (<https://www.r-project.org>). This study was approved by the Survey Behavioral Research Ethics Committee of The Chinese University of Hong Kong (reference no. SBRE-20-037).

Results

Study Timeline

The 5 study rounds intertwined epidemic waves 1 and 2 in Hong Kong (20) at different disease stages (Figure 1): the initial phase (R1), amid epidemic waves (R2 and R4), during the relative quiescence between 2 waves (R3), and the decaying phase of wave 2 (R5). The government-initiated interventions (such as school closure and compulsory mask-wearing) and the call for COVID-19 vaccine were also presented (Figure 1). The data collection was

completed before any announcement of the safety and efficacy trials of the candidate vaccines.

We received 2,478 attempts to complete the survey in R1, of which 1,715 provided complete responses (8), and 1,054 indicated willingness to participate in future studies. The sample sizes for R2–R5 ranged from 441 to 644 (Figure 2).

Respondent Characteristics

The cohort consisted of more females (68.5%–69.8%) and young adults (18–44 years of age) (78.6%–81.0%). Most were well educated: 78.9%–82.5% had at least postsecondary level education (Appendix Table 3). Most respondents were free from chronic diseases (87.1%–88.8%) and perceived themselves to be in good health (73.1%–78.1%) (Appendix Table 4).

Risk Perception over Time

We identified significant temporal variation of risk perception toward COVID-19 (Appendix Table 5). Assuming no precaution measures, respondents perceived themselves likely to be infected with COVID-19 (87.2%–92.8%). This proportion dropped to 19.3%–42.0% when the current situations were considered, when institutionalized interventions were in place and personal protective measures were conducted en masse (Appendix Table 1).

Perceived severity decreased significantly ($p < 0.001$) over the study period, from 97.4% (R1) of respondents considering COVID-19 to be serious to 77.2% (R5). The perceived chance of having COVID-19 cured increased significantly ($p < 0.001$) by more than 3-fold, from 16.6% (R1) to 57.2% (R5). An increasing time trend ($p < 0.001$) was also observed for perceived survival chance if infected, from 18.6% (R1) to 67.2% (R5).

Psychological Distress

The mean HADS-A score remained borderline abnormal throughout the study, ranging from 8.99 (R1) to 7.61 (R5). There was a substantial increase in the proportion of normal respondents in terms of anxiety ($p < 0.001$), from 35.6% (R1) to 51.7% (R5) (Appendix Table 6). This anxiety metric echoed the significant drop in the frequency of thinking about COVID-19 ($p < 0.001$), from 76.2% (R1) to 48.6% (R5). Despite this ease in anxiety level, the proportion of respondents worrying specifically about COVID-19 (85.7%–96.8%) and having their daily lives affected a lot by COVID-19 (45.7%–61.8%) remained high throughout the study (Appendix Table 6). Meanwhile, $\approx 40\%$ of the respondents showed symptoms of burnout toward COVID-19 since R3.

Adoption of Precautionary Measures

The adoption of individual precautionary measures remained high throughout the study (Appendix Table 1). Most respondents reported they wore masks (R1, 99.0%; R5, 99.8%), covered mouth and nose when coughing or sneezing (R1, 96.9%; R5, 98.4%), performed hand hygiene using hand sanitizer or alcohol gel (R1, 95.8%; R5, 97.7%), and disinfected their homes (R1, 78.6%; R5, 88.5%). The latter 2 measures showed a significant increasing trend across time.

The adoptions of social distancing across rounds were consistently from moderate to high (Appendix Table 1). About one third of the respondents avoided public transportation (R1, 38.0% to R5, 35.6%; with no significant trend, $p = 0.11$) and work (R1, 24.6% to R5, 35.4%; $p < 0.001$) across waves. Upward significant trends were observed among respondents in avoiding social activities (R1, 63.8% to R5, 85.7%; $p < 0.001$) and contacting with persons with fever or symptoms of respiratory diseases (R1, 92.9% to R5, 95.1%; $p < 0.05$).

Factors associated with greater adoption of social-distancing were being female (aOR 1.30, 95% CI 1.09–1.56), living in the New Territories, a suburb of Hong Kong, (aOR: 1.40–1.42), and being anxious (aOR 1.47, 95% CI 1.23–1.76) (Appendix Table 7). Respondents with chronic conditions (aOR 0.72, 95% CI 0.54–0.95) and those reporting having neutral understanding of COVID-19, compared with those who said they understood COVID-19 not well/not well at all (aOR 0.73, 95% CI 0.62–0.85) were less likely to practice social distancing (Appendix Table 7).

Likelihood of COVID-19 Vaccine Uptake

Respondents' intention to receive COVID-19 vaccine decreased significantly from R4 (48.7%, 95% CI 44.0–53.4) to R5 (37.6%, 95% CI 32.9–42.4), with particularly low rates among persons ≥ 55 years of age (Appendix Table 8). Factors associated with higher tendency for receiving COVID-19 vaccines were anxiety (borderline abnormal: aOR 1.53, 95% CI 1.04–2.23; abnormal: aOR 1.87, 95% CI 1.19–2.93), complacency (aOR 0.72, 95% CI 0.62–0.85), confidence (aOR 1.71, 95% CI 1.48–1.99), and collective responsibility (aOR 1.31, 95% CI 1.10–1.55). Compared with persons 18–24 years of age, persons ≥ 55 years of age were less likely to receive COVID-19 vaccine (aOR 0.47, 95% CI 0.23–0.98) (Appendix Table 8).

We also researched the trends of other psychobehavioral responses, including comparison of perceived severity across diseases (Appendix; Appendix Table 9), confidence in government and doctors (Appendix; Appendix Table 10), knowledge of COVID-19 (Appendix; Appendix

Table 11), and perceived effectiveness of precautionary measures (Appendix; Appendix Table 2).

Discussion

We conducted a 5-round longitudinal online survey to analyze the temporal changes in the community responses throughout the first 2 COVID-19 epidemics in Hong Kong. Overall, perceived susceptibility (assuming no precautionary measure taken) remained high: self-susceptibility (87.2%–92.8%) was substantially higher than that observed for the 2003 SARS (23.0%) (21) and the 2009 pandemic influenza (58.1%) (22) in the same population. However, in terms of perceived severity, the proportions dropped dramatically across time but were still higher than those observed in other highly affected countries (UK, 20.7% [12]; Kerala state, India, 55.7% [23]). The proportions of persons with an abnormal level of anxiety also dropped over the study period, from 34.3% to 22.0%. We observed consistently high levels of precautionary measures, such as mask wearing, hand hygiene, and home disinfection throughout the study period. Greater anxiety was associated with higher tendency of social distancing. The projected COVID-19 vaccine uptake rate dropped from 48.7% (R4) to 37.6% (R5). Greater anxiety, confidence in the vaccine, collective responsibility, and lower complacency contributed to a greater likelihood of intended vaccination.

Implications of Results

Our results have 5 immediate public health implications. First, with the uncertain disease progression (e.g., emergence of new variants of coronavirus) and the changing institutionalized interventions, there should be continual monitoring of risk perception toward COVID-19 in the community. Risk perception is an indispensable determinant of behavioral change (24) and depends on the prevalence of the health risk concerned (25). Our findings show time-varying risk perception amid pandemic times, with a perceived severity of COVID-19 that significantly decreased over time. Inferring from the large difference between naive (assuming no precautionary measures) and current (based on the current situation) scenarios, perceived susceptibility is sensitive to the disease progression and interventions in place. Although such temporal trend of risk perception was also observed in past pandemics (26), the absolute level of risk perception was not.

Second, surveillance and encouragement of social distancing should be maintained in the medium to long term, given the low projected uptake rate of COVID-19 vaccine. In Hong Kong, the reproductive number peaked at 2.39 in wave 1 and 3.04 in wave 2 (20), which (based on early data) corresponded approximately to requiring 56.1%–66.9% of the population to be immune to confer herd immunity (27). Because the projected vaccine uptake rates (R4, 48.7%; R5, 37.6%) fell short of the required level, relatively small-scale upcoming epidemics compared with the previous waves are expected. With more persons being vaccinated, there might be more social interactions. Effort should be allocated to maintain social distancing (such as avoiding unnecessary gatherings). Meanwhile, further research should focus on disease transmission during a mix of social distancing in place and vaccine hesitancy in the population.

Third, risk communications in Hong Kong should target complacency, vaccine confidence, and collective responsibility to boost the COVID-19 vaccine uptake. Temporal changes of the psychological antecedents of vaccine hesitancy should be closely monitored to inform the design of vaccination campaigns. We reported a low intention for uptake of the would-be vaccines, which declined over time in Hong Kong. A similar situation was observed in the United States, where the projected vaccine uptake rate dropped from 74.1% in April 2020 to 56.2% in December 2020 (28). Such low uptake intention among older persons in our study (R4, 29.4%; R5, 31.4%) is particularly worrisome because older age is a risk factor for death from COVID-19 (29).

The extent to which our findings on the predictors of uptake intention can be generalized to other countries or regions requires further investigation. Unique to Hong Kong were the low COVID-19 infection rate and low level of confidence in government measures. The weak uptake intention reported in this study was uncommon compared with other countries (overall 71.5%) (30). The low infection rate, along with the decreasing perceived severity toward COVID-19, might weaken the urgency for vaccination, which may also apply to places such as Taiwan, Japan, and Australia. However, the social unrest in Hong Kong in late 2019 might have led to distrust in the government (31), which could subsequently lower vaccination intention (32) and trigger maintenance of personal precautionary measures. One possible explanation is that, when moderated by distrust in government, persons tend to rely on personal protective measures (such as wearing facemasks and maintaining social distancing) but become skeptical to institutional protective measures (such as vaccines). Distrust in governments during the pandemic may also

influence vaccine hesitancy in other regions, such as Brazil and Poland (33). Nevertheless, given the projected low vaccine uptake rate in this study, it may be insufficient to reach herd immunity in the near future, if ever, in Hong Kong. Therefore, taking the vaccine or not may have little bearing on relaxing government interventions in the medium term. In addition, from findings in other regions, trust in the government itself (34) and the information provided by the government (30) increased preventive practices, specifically accepting vaccines, during pandemics (30). Therefore, effective health communication is particularly crucial for the Hong Kong government. To rebuild trust, public health authorities need to possess competence, objectivity, fairness, consistency, transparency, sincerity, and faith (35). In addition, modalities aside from government and healthcare providers, such as professional bodies and religious groups, may help deliver pro-vaccine messages (36).

Fourth, our results help to prioritize the content in promotional messaging. It is worth investing resources on promotional messaging, particularly when few respondents in R4 (overall, 16.7%; 18-24 years (y), 24.7%; 25-34y, 14.5%; 35-44y, 15.5%; 45-54y, 11.5%; $\geq 55y$, 17.6%) and R5 (overall, 10.5%; 18-24y, 12.8%; 25-34y, 7.4%; 35-44y, 12.1%; 45-54y, 6.1%; $\geq 55y$, 20.0%) indicated an absolute “yes” for receiving COVID-19 vaccines (measured on a 11-point Likert scale) and when there was antibody waning after receiving the vaccine. The decreasing “confidence” metric from R4 to R5 highlighted the need to build trust among the public toward the logistics of vaccine development, licensing, generating recommendations, and distribution (37). Before the government authorized the use of a COVID-19 vaccine, establishment of an advisory panel will help determine factors that the government should consider, such as performance (safety, efficacy, and effectiveness) and characteristics (number of doses, formulation, and presentation and packaging) of the available vaccine (38). Moreover, to increase the collective responsibility and perceived vaccine necessity, the government should foster understanding of the vaccine among the public with transparent communication, including more engagement with different stakeholders in the community and populations who are disproportionately affected by the pandemic to listen to their concerns. Leveraging knowledge, skills, and expertise from these communications will provide a robust assessment to underpin the vaccination campaign. Although calculations and constraints in the 5Cs model were not associated with the vaccine uptake likelihood at this stage, continuous examination in these 2 constructs will help refine future vaccination campaigns to engage citizens in cost–benefit calculations and increase their vaccine availability, affordability and accessibility.

Fifth, the psychological distress arising from burnout should be weighed together with the well-established anxiety. This COVID-19 epidemic is ongoing and has lasted much longer than the SARS epidemic, so more persons are developing syndromes of emotional exhaustion. The interplay between 2 psychological distresses, burnout and anxiety, is worth investigating during the ongoing epidemic. Our study showed that almost half of respondents had burnout symptoms in a short 4-month window from June to September 2020. This symptom did not contribute to the likelihood of COVID-19 vaccination in the last 2-point survey. However, the current general measure of burnout was not able to pinpoint the sources of burnout, such as financial stress, social isolation, the disease itself, or their combinations, for a detailed analysis. Nevertheless, the burnout phenomenon among persons coping with a long-term pandemic (39) suggests the need to reexamine the temporal association among social-distancing adoption, vaccination, and burnout.

This study has limitations. First, the survey may be subject to recall and social conformity biases, but its longitudinal design allowed us to track the same respondents over time, reducing self-control bias. Second, caution should be exercised when generalizing our findings to other regions because Hong Kong was exposed to other disease outbreaks recently, such as 1997 avian influenza (40), 2003 SARS (41), and the 2009 pandemic influenza (42). Nevertheless, our COVID-19 experience after those past outbreaks may be precedent to other countries, following the current COVID-19 pandemic. Third, our survey was conducted before the safety and efficacy data of the COVID-19 vaccines were released. The actual uptake rates might be affected by possible vaccination side effects events, such as the recent reported deaths after taking the vaccines in Hong Kong (43–45).

Continuous longitudinal assessment of community psychobehavioral responses during the pandemic can help public health authorities to tailor respective health communication strategies to achieve the desired behavioral outcomes (vaccination and adoption of precautionary measures) to control the upcoming waves of epidemic.

Acknowledgments

We thank Henry Chan and Sui Ting Fong for technical support. K.O. Kwok also acknowledges support from Health and Medical Research Fund (reference numbers: INF-CUHK-1, 17160302, 18170312), General Research Fund (reference numbers: 14112818, 24104920), Wellcome Trust Fund (United Kingdom, 200861/Z/16/Z), and Group Research Scheme of The Chinese University of Hong Kong.

About the Author

Dr. Kwok is an assistant professor affiliated with the JC School of Public Health and Primary Care at The Chinese University of Hong Kong, Hong Kong, China. His primary research interests include infectious disease epidemiology and infection control, with a current focus on mitigating emerging infectious disease outbreaks from the mathematic, epidemiologic, and behavioral perspectives.

References

- <eref>1. World Health Organization. COVID-19 [cited 2020 Dec 24].
<https://www.who.int/news/item/27-04-2020-who-timeline—covid-19></eref>
- <eref>2. World Health Organization. WHO coronavirus disease (COVID-19) dashboard; 2020 [cited 2020 Dec 24]. <https://covid19.who.int></eref>
- <jrn>3. Kwok KO, Cowling B, Wei V, Riley S, Read JM. Temporal variation of human encounters and the number of locations in which they occur: a longitudinal study of Hong Kong residents. *J R Soc Interface*. 2018;15:20170838. [PubMed <https://doi.org/10.1098/rsif.2017.0838>](https://doi.org/10.1098/rsif.2017.0838)</jrn>
- <jrn>4. Kwok KO, Wong V, Wei VWI, Wong SYS, Tang JW-T. Novel coronavirus (2019-nCoV) cases in Hong Kong and implications for further spread. *J Infect*. 2020;80:671–93. [PubMed <https://doi.org/10.1016/j.jinf.2020.02.002>](https://doi.org/10.1016/j.jinf.2020.02.002)</jrn>
- <jrn>5. Kwok KO, Huang Y, Tsoi MTF, Tang A, Wong SYS, Wei WI, et al. Epidemiology, clinical spectrum, viral kinetics and impact of COVID-19 in the Asia-Pacific region. *Respirology*. 2021;26:322–33. [PubMed <https://doi.org/10.1111/resp.14026>](https://doi.org/10.1111/resp.14026)</jrn>
- <eref>6. Centre for Health Protection. Countries/areas with reported cases of coronavirus disease-2019 (COVID-19); 2021 [cited 2021 Apr 9].
https://www.chp.gov.hk/files/pdf/statistics_of_the_cases_novel_coronavirus_infection_en.pdf</eref>
- <eref>7. World Health Organization. Draft landscape and tracker of COVID-19 candidate vaccines [cited 2020 Dec 24]. <https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines></eref>
- <jrn>8. Kwok KO, Li KK, Chan HHH, Yi YY, Tang A, Wei WI, et al. Community responses during early phase of COVID-19 epidemic, Hong Kong. *Emerg Infect Dis*. 2020;26:1575–9. [PubMed <https://doi.org/10.3201/eid2607.200500>](https://doi.org/10.3201/eid2607.200500)</jrn>

- <jrn>9. Stickley A, Matsubayashi T, Sueki H, Ueda M. COVID-19 preventive behaviours among people with anxiety and depressive symptoms: findings from Japan. *Public Health*. 2020;189:91–3. [PubMed https://doi.org/10.1016/j.puhe.2020.09.017](https://doi.org/10.1016/j.puhe.2020.09.017)</jrn>
- <jrn>10. Alanezi F, Aljahdali A, Alyousef S, Alrashed H, Alshaikh W, Mushcab H, et al. Implications of public understanding of COVID-19 in Saudi Arabia for fostering effective communication through awareness framework. *Front Public Health*. 2020;8:494. [PubMed https://doi.org/10.3389/fpubh.2020.00494](https://doi.org/10.3389/fpubh.2020.00494)</jrn>
- <jrn>11. Vai B, Cazzetta S, Ghiglini D, Parenti L, Saibene G, Toti M, et al. risk perception and media in shaping protective behaviors: insights from the early phase of COVID-19 Italian outbreak. *Front Psychol*. 2020;11:563426. [PubMed https://doi.org/10.3389/fpsyg.2020.563426](https://doi.org/10.3389/fpsyg.2020.563426)</jrn>
- <jrn>12. Bowman L, Kwok KO, Redd R, Yi Y, Ward H, Wei WI, et al. Comparing public perceptions and preventive behaviors during the early phase of the COVID-19 pandemic in Hong Kong and the United Kingdom: cross-sectional survey study. *J Med Internet Res*. 2021;23:e23231. [PubMed https://doi.org/10.2196/23231](https://doi.org/10.2196/23231)</jrn>
- <jrn>13. Coustasse A, Kimble C, Maxik K. COVID-19 and vaccine hesitancy: a challenge the United States must overcome. *J Ambul Care Manage*. 2021;44:71–5. [PubMed https://doi.org/10.1097/JAC.0000000000000360](https://doi.org/10.1097/JAC.0000000000000360)</jrn>
- <jrn>14. Ni MY, Yao XI, Leung KSM, Yau C, Leung CMC, Lun P, et al. Depression and post-traumatic stress during major social unrest in Hong Kong: a 10-year prospective cohort study. *Lancet*. 2020;395:273–84. [PubMed https://doi.org/10.1016/S0140-6736\(19\)33160-5](https://doi.org/10.1016/S0140-6736(19)33160-5)</jrn>
- <eref>15. Coalition for Epidemic Preparedness Innovations. \$2 billion required to develop a vaccine against the COVID-19 virus; 2020 [cited 2021 Jan 4]. https://cepi.net/news_cepi/2-billion-required-to-develop-a-vaccine-against-the-covid-19-virus-2/</eref>
- <jrn>16. Leung CM, Wing YK, Kwong PK, Lo A, Shum K. Validation of the Chinese-Cantonese version of the hospital anxiety and depression scale and comparison with the Hamilton Rating Scale of Depression. *Acta Psychiatr Scand*. 1999;100:456–61. [PubMed https://doi.org/10.1111/j.1600-0447.1999.tb10897.x](https://doi.org/10.1111/j.1600-0447.1999.tb10897.x)</jrn>
- <jrn>17. Dolan ED, Mohr D, Lempa M, Joos S, Fihn SD, Nelson KM, et al. Using a single item to measure burnout in primary care staff: a psychometric evaluation. *J Gen Intern Med*. 2015;30:582–7. [PubMed https://doi.org/10.1007/s11606-014-3112-6](https://doi.org/10.1007/s11606-014-3112-6)</jrn>

- <jrn>18. Betsch C, Schmid P, Heinemeier D, Korn L, Holtmann C, Böhm R. Beyond confidence: development of a measure assessing the 5C psychological antecedents of vaccination. *PLoS One*. 2018;13:e0208601. [PubMed https://doi.org/10.1371/journal.pone.0208601](https://doi.org/10.1371/journal.pone.0208601)</jrn>
- <eref>19. Derrick B, Dobson-Mckittrick A, Toher D, White P. Test statistics for comparing two proportions with partially overlapping samples; 2015 [cited 2021 Jan 4]. <https://uwe-repository.worktribe.com/output/805270></eref>
- <jrn>20. Kwok KO, Wei WI, Huang Y, Kam KM, Chan YYE, Riley S, et al. Evolving epidemiological characteristics of COVID-19 in Hong Kong, January to August 2020. *J Med Internet Res*. 2021 Mar 19 [Epub ahead of print]. [PubMed https://doi.org/10.2196/26645](https://doi.org/10.2196/26645)</jrn>
- <jrn>21. Leung GM, Quah S, Ho L-M, Ho S-Y, Hedley AJ, Lee H-P, et al. A tale of two cities: community psychobehavioral surveillance and related impact on outbreak control in Hong Kong and Singapore during the severe acute respiratory syndrome epidemic. *Infect Control Hosp Epidemiol*. 2004;25:1033–41. [PubMed https://doi.org/10.1086/502340](https://doi.org/10.1086/502340)</jrn>
- <jrn>22. Mak K-K, Lai C-M. Knowledge, risk perceptions, and preventive precautions among Hong Kong students during the 2009 influenza A (H1N1) pandemic. *Am J Infect Control*. 2012;40:273–5. [PubMed https://doi.org/10.1016/j.ajic.2011.10.023](https://doi.org/10.1016/j.ajic.2011.10.023)</jrn>
- <jrn>23. Jose R, Narendran M, Bindu A, Beevi N, L M, Benny PV. Public perception and preparedness for the pandemic COVID 19: A Health Belief Model approach. *Clin Epidemiol Glob Health*. 2021;9:41–6. [PubMed https://doi.org/10.1016/j.cegh.2020.06.009](https://doi.org/10.1016/j.cegh.2020.06.009)</jrn>
- <edb>24. Gaube S, Lermer E, Fischer P. The concept of risk perception in health-related behavior theory and behavior change. In: Raue M, Streicher B, Leimer E, editors. *Perceived Safety*. Springer International Publishing; 2019. p. 101–18.</edb>
- <jrn>25. El-Toukhy S. Parsing susceptibility and severity dimensions of health risk perceptions. *J Health Commun*. 2015;20:499–511. [PubMed https://doi.org/10.1080/10810730.2014.989342](https://doi.org/10.1080/10810730.2014.989342)</jrn>
- <jrn>26. Yeung NCY, Lau JTF, Choi KC, Griffiths S. Population responses during the pandemic phase of the influenza A(H1N1)pdm09 epidemic, Hong Kong, China. *Emerg Infect Dis*. 2017;23:813–5. [PubMed https://doi.org/10.3201/eid2305.160768](https://doi.org/10.3201/eid2305.160768)</jrn>
- <jrn>27. Kwok KO, Lai F, Wei WI, Wong SYS, Tang JWT. Herd immunity—estimating the level required to halt the COVID-19 epidemics in affected countries. *J Infect*. 2020;80:e32–3. [PubMed https://doi.org/10.1016/j.jinf.2020.03.027](https://doi.org/10.1016/j.jinf.2020.03.027)</jrn>

- <jrn>28. Szilagyi PG, Thomas K, Shah MD, Vizueta N, Cui Y, Vangala S, et al. National trends in the US public's likelihood of getting a COVID-19 vaccine—April 1 to December 8, 2020. *JAMA*. 2021;325:396–8. [PubMed https://doi.org/10.1001/jama.2020.26419](https://doi.org/10.1001/jama.2020.26419)</jrn>
- <jrn>29. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054–62. [PubMed https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)</jrn>
- <jrn>30. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A global survey of potential acceptance of a COVID-19 vaccine. *Nat Med*. . 2021;27:225–8. [PubMed https://doi.org/10.1038/s41591-020-1124-9](https://doi.org/10.1038/s41591-020-1124-9)</jrn>
- <jrn>31. Legido-Quigley H, Asgari N, Teo YY, Leung GM, Oshitani H, Fukuda K, et al. Are high-performing health systems resilient against the COVID-19 epidemic? *Lancet*. 2020;395:848–50. [PubMed https://doi.org/10.1016/S0140-6736\(20\)30551-1](https://doi.org/10.1016/S0140-6736(20)30551-1)</jrn>
- <jrn>32. Miyachi T, Takita M, Senoo Y, Yamamoto K. Lower trust in national government links to no history of vaccination. *Lancet*. 2020;395:31–2. [PubMed https://doi.org/10.1016/S0140-6736\(19\)32686-8](https://doi.org/10.1016/S0140-6736(19)32686-8)</jrn>
- <jrn>33. Lazarus JV, Ratzan S, Palayew A, Billari FC, Binagwaho A, Kimball S, et al. COVID-SCORE: A global survey to assess public perceptions of government responses to COVID-19 (COVID-SCORE-10). *PLoS One*. 2020;15:e0240011. [PubMed https://doi.org/10.1371/journal.pone.0240011](https://doi.org/10.1371/journal.pone.0240011)</jrn>
- <jrn>34. Min C, Shen F, Yu W, Chu Y. The relationship between government trust and preventive behaviors during the COVID-19 pandemic in China: exploring the roles of knowledge and negative emotion. *Prev Med*. 2020;141:106288. [PubMed https://doi.org/10.1016/j.ypmed.2020.106288](https://doi.org/10.1016/j.ypmed.2020.106288)</jrn>
- <eref>35. World Health Organization Europe. Vaccination and trust (2017); 2019 [cited 2021 Apr 7]. <https://www.euro.who.int/en/health-topics/disease-prevention/vaccines-and-immunization/publications/2017/vaccination-and-trust-2017></eref>
- <jrn>36. Lee C, Whetten K, Omer S, Pan W, Salmon D. Hurdles to herd immunity: distrust of government and vaccine refusal in the US, 2002–2003. *Vaccine*. 2016;34:3972–8. [PubMed https://doi.org/10.1016/j.vaccine.2016.06.048](https://doi.org/10.1016/j.vaccine.2016.06.048)</jrn>
- <jrn>37. Opel DJ, Salmon DA, Marcuse EK. Building trust to achieve confidence in COVID-19 vaccines. *JAMA Netw Open*. 2020;3:e2025672. [PubMed https://doi.org/10.1001/jamanetworkopen.2020.25672](https://doi.org/10.1001/jamanetworkopen.2020.25672)</jrn>

- <eref>38. World Health Organization. Principles and considerations for adding a vaccine to a national immunization programme; 2015 [cited 2021 Jan 1].
https://www.who.int/immunization/programmes_systems/policies_strategies/vaccine_intro_resources/nvi_guidelines/en</eref>
- <jrn>39. Carvalho Aguiar Melo M, de Sousa Soares D. Impact of social distancing on mental health during the COVID-19 pandemic: an urgent discussion. *Int J Soc Psychiatry*. 2020;66:625–6.
[PubMed <https://doi.org/10.1177/0020764020927047>](https://doi.org/10.1177/0020764020927047)</jrn>
- <jrn>40. Chan PKS. Outbreak of avian influenza A(H5N1) virus infection in Hong Kong in 1997. *Clin Infect Dis*. 2002;34(Suppl 2):S58–64. [PubMed <https://doi.org/10.1086/338820>](https://doi.org/10.1086/338820)</jrn>
- <jrn>41. Hung LS. The SARS epidemic in Hong Kong: what lessons have we learned? *J R Soc Med*. 2003;96:374–8. [PubMed <https://doi.org/10.1177/014107680309600803>](https://doi.org/10.1177/014107680309600803)</jrn>
- <jrn>42. Kwok KO, Davoudi B, Riley S, Pourbohloul B. Early real-time estimation of the basic reproduction number of emerging or reemerging infectious diseases in a community with heterogeneous contact pattern: using data from Hong Kong 2009 H1N1 pandemic influenza as an illustrative example. *PLoS One*. 2015;10:e0137959. [PubMed <https://doi.org/10.1371/journal.pone.0137959>](https://doi.org/10.1371/journal.pone.0137959)</jrn>
- <eref>43. Cheng S. Two more deaths, another nine temporary facial paralysis cases after Covid vaccinations; 2021 [cited 2021 Apr 10]. <https://hongkongfp.com/2021/03/25/two-more-deaths-another-nine-temporary-facial-paralysis-cases-after-covid-vaccinations></eref>
- <eref>44. Magramo K, Choy G, Tsang E. Coronavirus: chronically ill Hongkonger dies days after getting BioNTech Covid-19 vaccine, while experts push for return of international travel; 2021 [cited 2021 Apr 10]. <https://www.scmp.com/news/hong-kong/health-environment/article/3126158/coronavirus-tourism-lawmaker-urges-hong-kong></eref>
- <eref>45. The Standard. Another man died after receiving Sinovac vaccine; 2021 [cited 2021 Apr 10]. <https://www.thestandard.com.hk/breaking-news/section/4/168423/Another-man-died-after-receiving-Sinovac-vaccine></eref>

Address for correspondence: Kin On Kwok, Room 419, 4/F, JC School of Public Health and Primary Care Building, Prince of Wales Hospital, Shatin, New Territories, Hong Kong; email: kkokwok@cuhk.edu.hk; Wan In Wei, Room 427, 4/F, JC School of Public Health and Primary Care Building, Prince of Wales Hospital, Shatin, New Territories, Hong Kong; email: vivian1628@cuhk.edu.hk




Figure 1. Incidence and anxiety level by report date, and timeline of major interventions, Hong Kong, 2020. A, school closure: A1, school closure, Jan 25–May 26; A2, early start of summer holiday, Jul 13–Sep 22. B, government work-from-home arrangement: B1, Jan 29–Mar 1; B2, Mar 23–May 3; B3, Jul 20–Aug 23. C, ban on gatherings in public places: C1, group size limit 4 persons, Mar 29–May 7; C2, group size limit 8 persons, May 8–Jun 18; C3, group size limit 50 persons, Jun 19–Jul 14; C4, group size limit 4 persons, Jul 15–Jul 28; C5, group size limit 2 persons, Jul 29–Sep 10; C6, group size limit 4 persons, Sep 11–present. D, compulsory mask wearing: D1, on public transportation, Jul 15–present; D2, on public transportation and in public indoor areas, Jul 23–present; D3, on public transportation and in public indoor and outdoor areas, Jul 29–present (exemption for country parks or when engaging in strenuous physical activities in public outdoor spaces, Aug 28–present). E, regulations applied to restaurants, Mar 28–present: $\leq 50\%$ of premises capacity; tables ≥ 1.5 m apart; no more than 2, 4, or 8 persons per table; compulsory mask wearing except when consuming food or drink; compulsory body temperature screening before entry; hand sanitizer on premises; suspension of dine-in service for the following periods: E1, 6 pm–4:59 am, Jul 15–Jul 28; E2, at all times, Jul 29–30; E3, 6 pm–4:59 am, Jul 31–Aug 27; E4, 9 pm–4:59 am, Aug 28–Sep 3; E5, 12 am–4:59 am, Sep 18–present. F, business closures: F1, closures of bathhouses, party rooms, clubs, karaoke clubs, May 8–May 28; F2, closures (starting Jul 15)

of bathhouses, party rooms, clubs, karaoke clubs (all reopened Sep 17), swimming pools (suspended from Jul 29), sports premises (Jul 29–Aug 28), clubhouses (reopened Aug 28), beauty parlors (reopened Aug 28), massage establishments (reopened Sep 4), places of public entertainment (reopened Aug 28), places for amusement (reopened Sep 4), fitness centers (reopened Sep 4), and amusement game and mahjong-tin kau establishments (reopened Sep 11). G, COVID-19 vaccine development timeline: G1, WHO Convention of Global Research and Innovation, Feb 11–12; G2, WHO Global Research Roadmap prioritizing vaccine development, Jun 3; G3, draft landscape of candidate vaccines, Apr 11; G4, launch of COVID-19 Access Pool for sharing data for vaccine development, May 29; G5, funding commitment at Global Vaccine Summit for immunization in low-income countries, Jun 4; G6, call for US\$31.3 billion for therapeutics and vaccine development, Jun 26; G7, second summit on COVID-19 research and innovation, Jul 1–2; G8, engaging >150 countries in financing vaccines, Jul 15; G9, outline of global vaccine procurement, Aug 6; G10, WHO guidance on vaccine allocation between and within countries, Sep 14; G11, WHO calls for vaccine manufacturers to apply for prequalification, Oct 1.




Figure 2. Study timeline and participant recruitment. The numbers refer to the number of respondents who indicated willingness to participate in the respective survey round (in addition, for Round 1, their being aged ≥ 18 years, and their residence in Hong Kong for ≥ 5 days per week in the preceding month). They may or may not have completed the questionnaire.

Appendix. Additional information on psychobehavioral responses and likelihood of receiving COVID-19 vaccines during the pandemic, Hong Kong.